

***Morganjonesia* gen. nov. for two atypical *Corynespora* and *Teratosperma* species**

KAI ZHANG^{1*}, MIN QIAO², ZE-FEN YU²,
DE-WEI LI^{3,4}, RAFAEL F. CASTAÑEDA-RUIZ⁵

¹ Department of Landscaping, Shandong Yingcai University, Jinan, Shandong 250104, China

² Laboratory for Conservation and Utilization of Bio-resources,
Key Laboratory for Microbial Resources of the Ministry of Education,
Yunnan University, Kunming, Yunnan, 650091, China

³ The Connecticut Agricultural Experiment Station,
Valley Laboratory, 153 Cook Hill Road, Windsor, CT 06095, USA

⁴ Co-Innovation Center for Sustainable Forestry in Southern China,
Nanjing Forestry University, Nanjing, Jiangsu 210037, China

⁵ Instituto de Investigaciones Fundamentales en Agricultura Tropical Alejandro de Humboldt
(INIFAT), Calle 1 Esq. 2, Santiago de Las Vegas, C. Habana, Cuba, C.P. 17200

* CORRESPONDENCE TO: kaise0907@126.com

ABSTRACT—*Morganjonesia* is proposed as a new genus for two species, *Corynespora calophylli* and *Teratosperma litchii*, which have monotretic conidial ontogeny on the terminal conidiogenous cells and solitary, euseptate, verruculose, brown conidia with a single, subulate or acute, hyaline conidial appendage at the apex (sometimes with another downwardly-direct conidial appendage). Illustrations of both species are provided.

KEY WORDS—asexual *Ascomycota*, hyphomycetes, systematics

Introduction

Corynespora Güssow, typified by *C. mazei* Güssow [= *C. cassicola* (Berk. & M.A. Curtis) C.T. Wei], is distinguished by macronematous, mononematous, and generally unbranched, erect conidiophores and monotretic, terminal, integrated, determinate or with several doliiform, enteroblastic percurrent extension conidiogenous cells. The conidiogenous loci are apical, slightly

depressed, somewhat melanized, and thickened around the pore to produce solitary or blastocatenate, multidistoseptate, obclavate, cylindrical, obovate, oval, smooth-walled or verruculose, pale pigmented conidia usually with a slightly melanized basal scar (Morgan-Jones 1988, Seifert & al. 2011). *Corynespora calophylli* clearly deviates from the generic concept of *Corynespora*, because it is characterized by monotretic conidial ontogeny and solitary, 2-euseptate, obclavate to obpyriform, brown conidia with an apical, acute, hyaline conidial appendage, slightly papillate at the truncate base (Holubová-Jechová & Castañeda-Ruiz 1986).

Teratosperma Syd. & P. Syd., typified by *T. singulare* Syd & P. Syd., is characterized by macronematous, mononematous, generally unbranched, erect conidiophores and monoblastic, terminal, integrated, indeterminate conidiogenous cells with several annellidic enteroblastic percurrent extensions. The conidia are staurospore, tetrastrate, rostrate, brown to olivaceous brown, septate at the main axis, with 1–3-basal arms, and pale pigmented (Ellis 1971, Seifert & al. 2011). *Teratosperma litchii* clearly deviates from the generic concept of *Teratosperma*, because it is distinguished by a monotretic conidial ontogeny and solitary, 3(–4)-euseptate, obclavate, brown conidia with 1-2 apical, subulate, hyaline appendage that are slightly papillate at the truncate base (Matsushima 1980).

Here we propose a new genus, *Morganjonesia*, to accommodate these two atypical species.

Taxonomy

Morganjonesia R.F. Castañeda, K. Zhang & D.W. Li, **gen. nov.**

MB 830288

Differs from *Corynespora* species by its euseptate conidia with an acute or subulate, hyaline apical or downwardly direct conidial appendage; and differs from *Teratosperma* species by its monotretic conidial ontogeny.

TYPE SPECIES: *Corynespora calophylli* Hol.-Jech. & R.F. Castañeda [= *Morganjonesia calophylli* (Hol.-Jech. & R.F. Castañeda) R.F. Castañeda & al.]

ETYMOLOGY: Latin, *Morganjonesia*, in honor of Dr. Gareth Morgan-Jones in recognition of his contributions to the study of hyphomycetes.

CONIDIOPHORES macronematous, mononematous, unbranched, erect, straight, cylindrical, septate, smooth, brown to dark brown. CONIDIogenous CELLS monotretic, integrated, terminal, cylindrical, determinate or sometimes with one or several oblong enteroblastic percurrent extensions. Conidial secession schizolytic. CONIDIA solitary, acrogenous, obclavate to obpyriform, euseptate,




FIG. 1. *Morganjonesia calophylli* (holotype, INIFAT C84/150).

A. Conidia with single apical conidial appendages; B. Conidiogenous cells with tretic loci.




FIG. 2. *Morganjonesia litchii* (holotype, MFC 9048).
A. Conidia with two conidial appendages; B. Conidiogenous cells with tretic loci.

constricted at the septa, slightly papillate at the truncate base, brown or dark brown, smooth-walled or verruculose, with 1–2 acute or subulate, hyaline, smooth conidial appendages at the apex.

Morganjonesia calophylli (Hol.-Jech. & R.F. Castañeda.) R.F. Castañeda,
K. Zhang & D.W. Li, **comb. nov.** FIG. 1
MB 830290
≡ *Corynespora calophylli* Hol.-Jech. & R.F. Castañeda, Česká Mykol. 40(2): 83 (1986).

Morganjonesia litchii (Matsush.) K. Zhang & R.F. Castañeda, **comb. nov.** FIG. 2
MB 830291
≡ *Teratosperma litchii* Matsush., Matsush. Mycol. Mem. 1: 73 (1980).

NOTE: *Solicorynespora* R.F. Castañeda & W.B. Kendr., typified by *S. zapatensis* R.F. Castañeda & W.B. Kendr., is similar to *Morganjonesia* in the monotretic conidial ontogeny and the solitary, euseptate conidia (Hernández-Restrepo & al. 2014, Ma & al. 2016), but its protologue (Castañeda-Ruiz & Kendrick 1990) did not describe a hyaline, acute or subulate conidial appendage.

Acknowledgments

This work was financed by the National Natural Science Foundation Program of PR China (31870016). The authors express their sincere gratitude to Dr. Josiane S. Monteiro and Dr. Flavia Rodrigues Barbosa for their critical review of the manuscript. We acknowledge the facilities provided by Dr. P.M. Kirk and Dr. K. Bensch through the Index Fungorum and MycoBank websites, respectively. Dr. Lorelei Norvell's editorial review and Dr. Shaun Pennycook's nomenclature review are greatly appreciated.

Literature cited

- Castañeda-Ruiz RF, Kendrick B. 1990. Conidial fungi from Cuba: II. University of Waterloo Biology Series 33. 62 p.
- Ellis MB. 1971. Dematiaceous hyphomycetes. Kew, Commonwealth Mycological Institute.
- Hernández-Restrepo M, Castañeda-Ruiz RF, Gené J, Silvera-Simón C, Cano J, Guarro J. 2013. Two new species of *Solicorynespora* from Spain. Mycological Progress 13: 157–164. <https://doi.org/10.1007/s11557-013-0903-9>
- Holubová-Jechová V, Castañeda Ruiz RF. 1986. Studies on Hyphomycetes from Cuba III. New and interesting dematiaceous taxa from leaf litter. Česká Mykologie 40: 74–85.
- Ma J, Xia JW, Zhang XG, Luo YQ, Castañeda-Ruiz RF. 2016. *Solicorynespora* species associated with dead branches of subtropical forests in southern China. Cryptogamie, Mycologie 37: 37–44. <https://doi.org/10.7872/crym/v37.iss1.2016.37>
- Matsushima T. 1980. Saprophytic microfungi from Taiwan, part 1. Hyphomycetes. Matsushima Mycological Memoirs 1. 82 p.
- Morgan-Jones G. 1988. Notes on hyphomycetes. LX. *Corynespora matuszakii*, an undescribed species with narrow, cylindrical, catenate conidia and highly reduced conidial cell lumina. Mycotaxon 33: 483–487.
- Seifert K, Morgan-Jones G, Gams W, Kendrick B. 2011. The genera of hyphomycetes. CBS Biodiversity Series 9. 997 p.